

HARNACK INEQUALITIES FOR SYMMETRIC STABLE LÉVY PROCESSES

MARINA SERTIC

ABSTRACT. In this paper we consider Harnack inequalities with respect to a symmetric α -stable Lévy process X in $\mathbb{R}^d$, $\alpha \in (0, 2)$, $d \geq 2$. We study the example from the article [2]. There, the authors have associated the Harnack inequality with the relative Kato condition, which is a condition on the Lévy measure. By checking the condition, in the case $\alpha \in (0, 1)$, they have established that the Harnack inequality does not hold. We give an alternative proof of this fact, using the setting of [2]. We define the harmonic functions explicitly. For a given starting point of the process, we examine the probability of hitting a certain set at the first exit time of a unit ball.

Moreover, we also examine the weak Harnack inequality for a certain class of symmetric α -stable Lévy processes. We consider a symmetric α -stable Lévy process, $\alpha \in (0, 2)$, for which a spherical part μ of the Lévy measure is a spectral measure. In addition, we assume that μ is absolutely continuous with respect to the uniform measure σ on the sphere and impose certain bounds on the corresponding density. Eventually, we show that the weak Harnack inequality holds.

Acknowledgment. The author is thankful to Moritz Kassmann and Mateusz Kwaśnicki for valuable discussions.

1. INTRODUCTION

In the paper, we use the notation

$$B(x_0, r) = \{x : |x - x_0| \leq r\}, r \geq 0,$$

and

$$B_r = B(0, r), r \geq 0.$$

We consider a *symmetric α -stable Lévy process*, which has the characteristic function of the form

$$(1.1) \quad \mathbb{E}^0[e^{iu \cdot X_t}] = e^{-t\Phi(u)}, u \in \mathbb{R}^d, t \geq 0,$$

where the characteristic exponent Φ is given by

$$(1.2) \quad \Phi(u) = \int_{\mathbb{S}^{d-1}} |u \cdot \xi|^\alpha \mu(d\xi).$$

The author appreciates the support of the International Graduate College "Stochastics and Real World Models", Universität Bielefeld.

The measure μ is symmetric, finite and non-zero on $\mathbb{S}^{d-1}$ (see [9], Theorem 14.13). Let the measure μ be absolutely continuous with respect to the uniform surface measure on $\mathbb{S}^{d-1}$ and denote its density by f_μ .

The potential density (or the heat kernel) $p(t, x, y) = p(t, y - x)$ is determined by the Fourier transform

$$\int_{\mathbb{R}^d} e^{i\xi \cdot x} p(t, x) \, dx = e^{-t\Phi(\xi)}, \quad \xi \in \mathbb{R}^d, \quad t \geq 0.$$

Definition 1.1. The Green function (or the potential kernel) is defined by

$$(1.3) \quad G(x, y) = \int_0^\infty p(t, x, y) \, dt, \quad x, y \in \mathbb{R}^d.$$

Definition 1.2. Let D be an open set, $D \subset \mathbb{R}^d$. The Green function of X^D is defined by

$$(1.4) \quad G_D(x, y) = G(x, y) - \mathbb{E}^x[G(X_{\tau_D}, y)], \quad x, y \in D.$$

2. HARNACK INEQUALITY

Definition 2.1. A non-negative Borel measurable function $u : \mathbb{R}^d \rightarrow \mathbb{R}$ is *harmonic* in an open set $D \subset \mathbb{R}^d$ with respect to the process X if

$$u(x) = \mathbb{E}^x[u(X_{\tau_U})], \quad x \in U,$$

for every bounded open set U such that $\bar{U} \subset D$. Here, $\tau_U = \inf\{t > 0 : X_t \notin U\}$ is the first exit time from the set U .

Definition 2.2. The Harnack inequality for a symmetric α -stable Lévy processes X holds true if there is a constant $K \geq 1$ such that for every non-negative function $u : \mathbb{R}^d \rightarrow \mathbb{R}$ which is harmonic in B_1 with respect to X , the inequality

$$u(x) \leq K \cdot u(y), \quad x, y \in B_{1/2}.$$

holds true.

2.1. Construction of the Sequence of Harmonic Functions.

Lemma 2.3. Let $c > 1$. There are sequences (α_n) and (β_n) of positive numbers such that:

$$(2.1) \quad A = \sum_{n=1}^\infty \alpha_n < \frac{\pi}{2}, \quad B = \sum_{n=1}^\infty \beta_n < \frac{\pi}{2},$$

$$(2.2) \quad \lim_{n \rightarrow \infty} (A_n + B_n) = A + B = \frac{\pi}{2},$$

$$(2.3) \quad \lim_{n \rightarrow \infty} \frac{\tan(A_n + B_{n+1})}{\tan(A_n + B_n)} = +\infty,$$

$$(2.4) \quad (c+2) \cdot \tan(A_n + B_{n+1}) \leq (c-1) \cdot \tan(A_{n+1} + B_{n+1}), \quad n \in \mathbb{N},$$

where A_n and B_n denote the partial sum of the sequences (α_n) and (β_n) , respectively, i.e. $A_n = \sum_{k=1}^n \alpha_k$, $B_n = \sum_{k=1}^n \beta_k$.

Proof. We construct inductively sequences (α_n) and (β_n) of positive numbers which fulfill the conditions of the lemma. Set $K = (c + 2)/(c - 1)$.

Choose α_1 and β_1 so that

$$\alpha_1 + \beta_1 < \pi/2.$$

Next, choose β_2 and α_2 so that:

$$\begin{aligned} \beta_2 &< \pi/2 - (A_1 + B_1), \\ \arctan(K \cdot \tan(A_1 + B_2)) - (A_1 + B_2) &< \alpha_2 < \pi/2 - (A_1 + B_2). \end{aligned}$$

Assume $\beta_1, \dots, \beta_n, \alpha_1, \dots, \alpha_n$ are chosen. Choose β_{n+1} and α_{n+1} so that:

$$\begin{aligned} \beta_{n+1} &< \pi/2 - (A_n + B_n), \\ n + 1 &\leq \frac{\tan(A_n + B_{n+1})}{\tan(A_n + B_n)}, \\ \arctan(K \cdot \tan(A_n + B_{n+1})) - (A_n + B_{n+1}) &< \alpha_{n+1} \\ \alpha_{n+1} &< \pi/2 - (A_n + B_{n+1}). \end{aligned}$$

Notice that the choice of the sequence (β_n) is possible, since

$$\lim_{h \nearrow \pi/2 - c} \frac{\tan(h + c)}{\tan(c)} = +\infty,$$

for fixed $c \in (0, \frac{\pi}{2})$.

Furthermore, the choice of the sequence (α_n) is possible due to the choice of (β_n) .

Now, from the choice of (α_n) , we have

$$\arctan(K \cdot \tan(A_n + B_{n+1})) < A_{n+1} + B_{n+1} < \pi/2,$$

which, together with the choice of (β_n) , implies

$$\lim_{n \rightarrow \infty} (A_n + B_n) = \frac{\pi}{2}.$$

□

Remark 2.4. At this point, we would like to mention that none of the following sequences satisfies the property (2.3) from the Lemma 2.3.

- (1) $\alpha_k = 2^{-k}$, $\beta_k = \frac{1}{k(k+1)}$
- (2) $\alpha_k = a^{-k}$, $\beta_k = b^{-k}$, $1 < b \leq a$,
- (3) $\alpha_k = \frac{1}{k(k+1)}$, $\beta_k = k^{-1-\delta}$, $\delta > 0$,
- (4) $\alpha_k = 2^{-(k^3)}$, $\beta_k = 2^{-(k^2)}$,
 $\alpha_k = a^{-(k^3)}$, $\beta_k = b^{-(k^2)}$, $a > 1, b > 1$

The details can be found in Lemma 4.1 (Appendix 4).



Definition 2.5. Let $\delta, x \in (0, 1)$ and (α_n) and (β_n) be sequences as in Lemma 2.3. We define the sets

$$\begin{aligned} S_n^\delta(x) &= \{(\delta, y) : A_{n-1} + B_n < \arctan\left(\frac{y}{x + \delta}\right) < A_n + B_n\}, \\ L_n^\delta(x) &= \bigcup_{z \in [x, 1]} S_n^\delta(z). \end{aligned}$$

Remark 2.6. Notice that, by the definitions of $S_n^\delta(x)$ and $L_n^\delta(x)$, we have

$$(2.5) \quad L_n^\delta(x) = \{(\delta, y) : (x + \delta) \cdot \tan(A_{n-1} + B_n) < y < (1 + \delta) \tan(A_n + B_n)\}.$$

The sets $S_n^\delta(x)$ and $L_n^\delta(x)$ are illustrated at the pictures that follow.



Remark 2.7. $S_n^\delta(x)$ describes the set on the line $\{(x, y) : x = \delta, y > 0\}$, which is seen from the point $(-x, 0)$ by the cone K_n , and $L_n^\delta(x)$ describes the set on the line $\{(x, y) : x = \delta, y > 0\}$ which is seen from the set $[-1, -x] \times \{0\}$ by the cone K_n .

Proposition 2.1. *Let (α_n) and (β_n) be sequences as in Lemma 2.3. For every $\delta \in (0, 1)$ and every $x \in (0, 1)$ there is $n_0 = n_0(\delta, x)$ such that for every $n \geq n_0$*

$$L_n^\delta(x) \cap L_{n+1}^\delta(x) = \emptyset.$$

Proof. Let $\delta, x \in (0, 1)$. Recall, by (2.5), we have

$$L_n^\delta(x) = \{(\delta, y) : (x + \delta) \cdot \tan(A_{n-1} + B_n) < y < (1 + \delta) \tan(A_n + B_n)\}.$$

Now, from Lemma 2.3, condition (2.3), it follows that there is $n_0 = n_0(\delta, x)$ such that for every $n \geq n_0$,

$$L_n^\delta(x) \cap L_{n+1}^\delta(x) = \emptyset,$$

which proves the proposition. $\square$

Definition 2.8. Let $\delta \in (0, 1)$ and (α_n) and (β_n) be sequences as in Lemma 2.3. For $z = (z_1, z_2), y = (y_1, y_2) \in \mathbb{R}^2$, we define the sets

$$\begin{aligned} \tilde{S}_n^\delta(z) &= \{(\delta, u) : A_{n-1} + B_n < \arctan\left(\frac{u - z_2}{\delta + z_1}\right) < A_n + B_n\}, \\ \tilde{L}_n^\delta(y) &= \bigcup_{\{z : z_1 \in [y_1, 1], z_2 = y_2\}} \tilde{S}_n^\delta(z). \end{aligned}$$


Remark 2.9. $\tilde{S}_n^\delta(z_1, z_2)$ describes the set on the line $\{(x, u) : x = \delta, u > 0\}$, which is seen from the point $(-z_1, z_2)$ by the cone K_n , and $\tilde{L}_n^\delta(y_1, y_2)$ describes the set on the line $\{(x, u) : x = \delta, u > 0\}$, which is seen from the set $[-1, -y_1] \times \{y_2\}$ by the cone K_n .

Remark 2.10. Let $w = (w_1, w_2) \in \mathbb{R}^2$. Notice that by the definitions of $\tilde{S}_n^\delta(z)$ and $\tilde{L}_n^\delta(y)$, we have

$$(2.6) \quad \begin{aligned} \tilde{L}_n^\delta(w) &= \{(\delta, u) : (w_1 + \delta) \cdot \tan(A_{n-1} + B_n) + w_2 < \\ &< u < (1 + \delta) \cdot \tan(A_n + B_n) + w_2\}. \end{aligned}$$

By means of the two propositions below, we will define the sequence of harmonic functions so that the Harnack inequality does not hold. To be more precise, we will construct the sequence of sets (B_n) and consequently the sequence of functions (u_n) , so that

$$\lim_{n \rightarrow \infty} \frac{u_n(1/2, 0)}{u_n(0, 0)} = 0.$$



Proposition 2.2. *Let (α_n) , (β_n) be sequences as in Lemma 2.3. There are sequences (δ_n) , (x_n) , (y_n) , (c_n) and (d_n) of positive numbers such that*

$$V_n \subset \tilde{S}_{n+1}^{\delta_n}(-x_1, x_2) \cap \Delta_n^{\delta_n},$$

for every $x = (x_1, x_2) \in U_n$, where

$$\begin{aligned} U_n &:= (-y_n, y_n)^2, \\ V_n &:= \{\delta_n\} \times (c_n, d_n). \end{aligned}$$

Remark 2.11. Notice that the picture above is in the connection with the statement.

Proof. For $n \in \mathbb{N}$, define

$$(2.7) \quad I_n := 2 + \tan(A_n + B_{n+1}) + \tan(A_{n+1} + B_{n+1}),$$

$$(2.8) \quad P_n := \tan(A_{n+1} + B_{n+1}) - \tan(A_n + B_{n+1}),$$

$$(2.9) \quad P'_n := \tan(A_n + B_{n+1}) - \tan(A_n + B_n).$$

Furthermore, set

$$(2.10) \quad \delta_n := \frac{I_n \cdot (1 + \tan(A_n + B_n)) \cdot (1 + \tan(A_{n+1} + B_{n+1}))}{P_n^2 \cdot (1 + \tan(A_n + B_{n+1})) + P_n' I_n \cdot (1 + \tan(A_{n+1} + B_{n+1}))},$$

$$(2.11) \quad x_n := 2\delta_n \cdot \frac{1 + \tan(A_n + B_{n+1})}{\tan(A_n + B_{n+1})} \cdot \frac{P_n}{I_n} + \frac{1}{\tan(A_n + B_{n+1})},$$

$$(2.12) \quad y_n := \frac{1 + \tan(A_n + B_n) - \delta_n \cdot P_n'}{1 + \tan(A_n + B_{n+1})},$$

$$(2.13) \quad c_n := (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n,$$

$$(2.14) \quad d_n := (\delta_n - y_n) \cdot \tan(A_{n+1} + B_{n+1}) - y_n.$$

For the convenience, let us denote the denominator of δ_n by H_n ,

$$(2.15) \quad H_n := P_n^2 \cdot (1 + \tan(A_n + B_{n+1})) + P_n' I_n \cdot (1 + \tan(A_{n+1} + B_{n+1})).$$

First, from Lemma 4.4 (Appendix 4), we have that I_n , P_n , P_n' , δ_n , x_n , y_n , c_n and d_n are positive and $c_n < d_n$.

Let us prove $V_n \subset \Delta_n^{\delta_n}$. What is more, we show $V_n = \Delta_n^{\delta_n}$. Recall, we defined the set $\Delta_n^{\delta_n}$ in the way

$$\Delta_n^{\delta_n} = \{(y_1, y_2) : y_1 = \delta_n, (\delta_n + 1) \cdot \tan(A_n + B_n) + 1 < y_2 < (\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1\}.$$

The definition (2.12) yields

$$(\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n = (\delta_n + 1) \cdot \tan(A_n + B_n) + 1.$$

Therefore, using (2.13) the equality

$$c_n = (\delta_n + 1) \cdot \tan(A_n + B_n) + 1$$

holds.

By Lemma 4.4 (Appendix 4) we have $d_n = (\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1$, hence $V_n = \Delta_n^{\delta_n}$.

Now, for $x = (x_1, x_2) \in U_n$, we prove $V_n \subset \tilde{S}_{n+1}^{\delta_n}(-x_1, x_2)$. Because of geometrical reasoning, it is enough to show

$$V_n \subset \tilde{S}_{n+1}^{\delta_n}(-y_n, -y_n) \cap \tilde{S}_{n+1}^{\delta_n}(y_n, y_n).$$

Let p_1^n and p_2^n be the lines given by

$$\begin{aligned} y - y_n &= \tan(A_n + B_{n+1}) \cdot (x + y_n), \\ y + y_n &= \tan(A_{n+1} + B_{n+1}) \cdot (x - y_n), \end{aligned}$$

respectively. As in the picture, p_1^n and p_2^n are the lines through the points $(-y_n, y_n)$ and $(y_n, -y_n)$, respectively.

Denote by P_1^n and P_2^n the points which determine the intersection of lines p_1^n and p_2^n with the set $\{(y_1, y_2) : y_1 = \delta_n, y_2 > 0\}$:


$$\begin{aligned} P_1^n &= (x_{P_1}, y_{P_1}) = (\delta_n, (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n), \\ P_2^n &= (x_{P_2}, y_{P_2}) = (\delta_n, (\delta_n - y_n) \cdot \tan(A_{n+1} + B_{n+1}) - y_n). \end{aligned}$$

Notice that P_1^n and P_2^n determine the lower boundary point and the upper boundary point of the sets $\tilde{S}_{n+1}^{\delta_n}(y_n, y_n)$ and $\tilde{S}_{n+1}^{\delta_n}(-y_n, -y_n)$, respectively. Notice

that showing $V_n = \{\delta_n\} \times (y_{P_1}, y_{P_2})$ finishes the proof. Clearly, the definitions (2.13) and (2.14) yield

$$\begin{aligned} c_n &= y_{P_1}, \\ d_n &= y_{P_2}, \end{aligned}$$

and hence the proposition. $\square$



Proposition 2.3. *Let (α_n) , (β_n) be sequences as in the Lemma 2.3 and let (I_n) , (P_n) , (P'_n) , (δ_n) , (x_n) , (y_n) , (c_n) , (d_n) and (U_n) be as in the Proposition 2.2. There exist sequences (δ'_n) and $(B(S_n, r_n))$, where:*

- (1) (δ'_n) is such that $\delta'_n < \delta_n$, for every $n \in \mathbb{N}$,
- (2) $B(S_n, r_n)$ is a ball with the center $S_n = (x_{S_n}, y_{S_n})$ and radius r_n , such that

$$B(S_n, r_n) \subset \bigcup_{\{\varepsilon: \delta'_n < \varepsilon < \delta_n\}} \left(\tilde{S}_{n+1}^\varepsilon(-x_1, x_2) \cap \Delta_n^\varepsilon \right),$$

for every $x = (x_1, x_2) \in U_n$.

Remark 2.12. Notions from the Proposition are illustrated at the picture above.

Proof. For $n \in \mathbb{N}$, define

$$(2.16) \quad \delta'_n := \frac{\delta_n \cdot (1 + \tan(A_n + B_{n+1})) + y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})},$$

$$(2.17) \quad \begin{aligned} J_n &:= 2 \tan(A_n + B_{n+1}) \cdot \tan(A_{n+1} + B_{n+1}) \\ &\quad + \tan(A_n + B_{n+1}) + \tan(A_{n+1} + B_{n+1}). \end{aligned}$$

Set

$$(2.18) \quad x_{S_n} := \frac{\delta_n + y_n}{2} \cdot \frac{I_n}{1 + \tan(A_{n+1} + B_{n+1})},$$

$$(2.19) \quad y_{S_n} := \frac{\delta_n + y_n}{2} \cdot \frac{J_n}{1 + \tan(A_{n+1} + B_{n+1})},$$

$$(2.20) \quad r_n := \frac{1}{2} \cdot \frac{\delta_n P_n - y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})}.$$

By the first part of Lemma 4.5 (Appendix 4),

$$B(S_n, r_n) \subset Q_n := (\delta'_n, \delta_n) \times (c_n, c_n + (\delta_n - \delta'_n)).$$

Now, it is enough to show that for every $x = (x_1, x_2) \in U_n$

$$Q_n \subset \bigcup_{\{\varepsilon: \delta'_n < \varepsilon < \delta_n\}} \left(\tilde{S}_{n+1}^\varepsilon(-x_1, x_2) \cap \Delta_n^\varepsilon \right).$$

Let $z = (z_1, z_2) \in Q_n$. We show

$$z \in \tilde{S}_{n+1}^\varepsilon(-x_1, x_2) \cap \Delta_n^\varepsilon,$$

for some $\delta'_n < \varepsilon < \delta_n$.

Recall,

$$\begin{aligned} \Delta_n^\varepsilon &= \{(y_1, y_2) : y_1 = \varepsilon, (\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 < \\ &\quad < y_2 < (\varepsilon + x_n) \cdot \tan(A_n + B_{n+1}) - 1\}, \end{aligned}$$

so in order to show $z \in \Delta_n^\varepsilon$, we prove that

$$\begin{aligned} c_n &\geq (\varepsilon + 1) \cdot \tan(A_n + B_n) + 1, \\ c_n + (\delta_n - \delta'_n) &\leq (\varepsilon + x_n) \cdot \tan(A_n + B_{n+1}) - 1, \end{aligned}$$

for some $\delta'_n < \varepsilon < \delta_n$. Since the proof is quite technical, we spell out the details in the second part of Lemma 4.5 (Appendix 4).

Now, for $x = (x_1, x_2) \in U_n$, let us prove $z \in \tilde{S}_{n+1}^\varepsilon(-x_1, x_2)$. Due to geometrical reasoning, it is enough to show that

$$z \in \tilde{S}_{n+1}^\varepsilon(-y_n, -y_n) \cap \tilde{S}_{n+1}^\varepsilon(y_n, y_n).$$

As before, let p_1^n and p_2^n denote the lines

$$\begin{aligned} y - y_n &= \tan(A_n + B_{n+1}) \cdot (x + y_n), \\ y + y_n &= \tan(A_{n+1} + B_{n+1}) \cdot (x - y_n), \end{aligned}$$

respectively. As before, the lines p_1^n and p_2^n are the lines through the points $(-y_n, y_n)$ and $(y_n, -y_n)$, respectively.

Denote by P_1^n and P_2^n the points which determine the intersection of p_1^n and p_2^n with the set $\{(y_1, y_2) : y_1 = \varepsilon, y_2 > 0\}$:

$$(2.21) \quad P_1^n = (x_{P_1}, y_{P_1}) = (\varepsilon, (\varepsilon + y_n) \cdot \tan(A_n + B_{n+1}) + y_n),$$

$$(2.22) \quad P_2^n = (x_{P_2}, y_{P_2}) = (\varepsilon, (\varepsilon - y_n) \cdot \tan(A_{n+1} + B_{n+1}) - y_n).$$


Notice that P_1^n and P_2^n determine the lower boundary point and the upper boundary point of sets $\tilde{S}_{n+1}^\varepsilon(y_n, y_n)$ and $\tilde{S}_{n+1}^\varepsilon(-y_n, -y_n)$, respectively. If we show $z_2 \in (y_{P_1}, y_{P_2})$, the proposition is proved.

To this end, we prove

$$\begin{aligned} c_n &\geq y_{P_1}, \\ c_n + (\delta_n - \delta'_n) &\leq y_{P_2}, \end{aligned}$$

and the details can be found in the Lemma 4.6 in Appendix 4. $\square$

2.2. Harnack Inequality. In the subsection that follows, we give a proof of the fact that Harnack inequality does not hold.



Let X be a symmetric α -stable Lévy process in $\mathbb{R}^2$, $0 < \alpha < 1$, with the characteristic function of the form

$$\mathbb{E}^0 e^{iu \cdot X_t} = e^{-t\Phi(u)}, \quad u \in \mathbb{R}^2,$$

where

$$\Phi(u) = \int_{\mathbb{S}^1} |u \cdot \xi|^\alpha \mu(d\xi).$$

We define the spectral measure μ appropriately, below in Theorem 2.14. Furthermore, for a given starting point $x \in B_1$ of the process X , we examine the probability

$$\mathbb{P}^x(X_{\tau_{B_1}} \in B_n), \quad n \in \mathbb{N}$$

of hitting the set $B_n = B(S_n, r_n)$ at the first exit time of a unit ball B_1 , where for every $n \in \mathbb{N}$, B_n are as in Proposition 2.3. We choose the points $w_0 = (1/2, 0)$, $0 = (0, 0)$ and define the harmonic functions

$$u_n(x) = \mathbb{P}^x(X_{\tau_{B_1}} \in B_n), \quad n \in \mathbb{N}, \quad x \in B_1.$$

We show

$$\frac{u_n(w_0)}{u_n(0)} \leq c(\alpha) \cdot a_n b_n,$$

for sequences (a_n) and (b_n) with certain properties and eventually obtain the desired conclusion.

Remark 2.13. Constants are positive real numbers, which exact value may vary from one line to the other. For the convenience they will not be explicitly stated throughout the proof.

Before we prove the Theorem 2.14, we state the proposition for the Green function estimate of the unit ball B_1 for X (see [2], Theorem 2).

Proposition 2.4. *Let X be a symmetric α -stable Lévy process in $\mathbb{R}^d$, $0 < \alpha < 2$, with the characteristic function of the form*

$$\mathbb{E}^0 e^{iu \cdot X_t} = e^{-t\Phi(u)}, \quad u \in \mathbb{R}^d,$$

where

$$\Phi(u) = \int_{\mathbb{S}^{d-1}} |u \cdot \xi|^\alpha \mu(d\xi),$$

and μ is a finite, symmetric measure on $\mathbb{S}^{d-1}$ such that $\mu(\mathbb{S}^{d-1}) > 0$. Let μ be absolutely continuous with respect to the surface measure on $\mathbb{S}^{d-1}$, and denote by f_μ its density, for which the inequality

$$0 \leq f_\mu(\xi) \leq m, \quad \xi \in \mathbb{S}^{d-1},$$

holds, for some $m > 0$. Then the estimate for the Green function G_{B_1}

$$c^{-1} \cdot s(y) \cdot |x - y|^{\alpha-d} \leq G_{B_1}(x, y) \leq c \cdot s(y) \cdot |x - y|^{\alpha-d},$$

holds, for all $|x| < 1/2, |y| < 1$. Here, $s(y) = \mathbb{E}^y[\tau_{B_1}]$ is the expected time spent in B_1 , if the process starts from $y \in B_1$.

Theorem 2.14. *Let X be a symmetric α -stable Lévy process in $\mathbb{R}^2$, $0 < \alpha < 1$, with the characteristic function of the form*

$$\mathbb{E}^0 e^{iu \cdot X_t} = e^{-t\Phi(u)}, \quad u \in \mathbb{R}^2,$$

where

$$\Phi(u) = \int_{\mathbb{S}^1} |u \cdot \xi|^\alpha \mu(d\xi).$$

Let the measure μ on $\mathbb{S}^1$ be absolutely continuous with respect to the surface measure on $\mathbb{S}^1$, and denote by f_μ its density. Set

$$f_\mu = \mathbf{1}_{\bigcup_{n \geq 1} (B_{\xi_n, r_n} \cup B_{-\xi_n, r_n})},$$

where

$$\begin{aligned} B_{\xi_n, r_n} &= B(\xi_n, r_n) \cap \mathbb{S}^1, \\ r_n &= 2 \cdot \sin(\alpha_n/4), \\ \xi_n &= (\cos(A_{n-1} + B_n + \alpha_n/2), \sin(A_{n-1} + B_n + \alpha_n/2)) \in \mathbb{S}^1, n \in \mathbb{N}, \end{aligned}$$

where (α_n) and (β_n) are as in Lemma 2.3. Then the Harnack inequality for X does not hold.

Proof. Let

$$\begin{aligned} u_n(x) &= \mathbb{P}^x(X_{\tau_B} \in B_n), \quad n \in \mathbb{N}, \quad x \in B(0, 1), \\ O_n &= ((-x_n, \delta_n) \times (-1, 1)) \cap B_1, \end{aligned}$$

where $B_n = B(S_n, r_n)$, x_n, δ_n are introduced in the Propositions 2.2 and 2.3, and set $w_0 = (1/2, 0)$ and $0 = (0, 0)$.

By the strong Markov property, every function u_n is harmonic in B_1 with respect to X .

Namely, for a set $U \subset \bar{U} \subset B_1$,

$$\begin{aligned} \mathbb{E}^x[u_n(X_{\tau_U})] &= \mathbb{E}^x[\mathbb{P}^{X_{\tau_U}}[X_{\tau_B} \in B_n]] \\ &= \mathbb{P}^x[X_{\tau_B} \circ \theta_{\tau_U} \in B_n] \\ &= \mathbb{P}^x[X_{\tau_B} \in B_n] \\ &= u_n(x). \end{aligned}$$

Using the Lévy system formula ([4]) it follows

$$\begin{aligned} &\mathbb{E}^{w_0} \left[\sum_{s \leq t \wedge \tau_B} \mathbf{1}_{\{X_{s-} \in O_n, X_s \in B_n\}} \right] \\ &= \mathbb{E}^{w_0} \left[\int_0^{t \wedge \tau_B} \mathbf{1}_{O_n}(X_s) \left(\int_{B_n} f_\nu(y - X_s) \, dy \right) \, ds \right] \\ &\leq \mathbb{E}^{w_0} \left[\int_0^{t \wedge \tau_B} \mathbf{1}_{O_n}(X_s) \left(\int_{B_n} |y - X_s|^{-\alpha-2} \, dy \right) \, ds \right] \\ &\leq |B_n| \cdot [(\delta_n + 1) \cdot \tan(A_n + B_n)]^{-\alpha-2} \cdot \mathbb{E}^{w_0} \left[\int_0^{t \wedge \tau_B} \mathbf{1}_{O_n}(X_s) \, ds \right] \\ &\leq |B_n| \cdot [(\delta_n + 1) \cdot \tan(A_n + B_n)]^{-\alpha-2} \cdot \mathbb{E}^{w_0} \left[\int_0^\infty \mathbf{1}_{O_n}(X_s) \, ds \right]. \end{aligned}$$

The estimates of the potential kernel ([10]) imply

$$\begin{aligned} \mathbb{E}^{w_0} \left[\int_0^\infty \mathbf{1}_{O_n}(X_s) \, ds \right] &\leq c(\alpha) \cdot \int_{O_n} |y - w_0|^{\alpha-2} \, dy \\ &\leq c(\alpha) \cdot (\delta_n + x_n). \end{aligned}$$

From here we have

$$\begin{aligned} & \mathbb{E}^{w_0} \left[\sum_{s \leq t \wedge \tau_B} \mathbf{1}_{\{X_{s-} \in O_n, X_s \in B_n\}} \right] \\ & \leq c(\alpha) \cdot |B_n| \cdot [(\delta_n + 1) \cdot \tan(A_n + B_n)]^{-\alpha-2} \cdot (\delta_n + x_n). \end{aligned}$$

Letting $t \rightarrow \infty$, by the dominated convergence theorem, we obtain

$$(2.23) \quad u_n(w_0) \leq c_1(\alpha, n) \cdot (\delta_n + x_n),$$

where

$$c_1(\alpha, n) = c(\alpha) \cdot |B_n| \cdot [(\delta_n + 1) \cdot \tan(A_n + B_n)]^{-\alpha-2}.$$

Let us compute the lower bound on $u_n(0)$.

By the Lévy system formula (see [4]) and the construction, it follows that

$$\begin{aligned} u_n(0) &= \mathbb{P}^0(X_{\tau_B} \in B_n) \\ &\geq \mathbb{E}^0 \left[\sum_{s \leq t \wedge \tau_B} \mathbf{1}_{\{X_{s-} \in U_n, X_s \in B_n\}} \right] \\ &= \mathbb{E}^0 \left[\int_0^{t \wedge \tau_B} \mathbf{1}_{U_n}(X_s) \left(\int_{B_n} f_\nu(y - X_s) \, dy \right) \, ds \right] \\ &= \mathbb{E}^0 \left[\int_0^{t \wedge \tau_B} \mathbf{1}_{U_n}(X_s) \left(\int_{B_n} |y - X_s|^{-\alpha-2} \, dy \right) \, ds \right] \\ &\geq \bar{c}_2(\alpha, n) \cdot \mathbb{E}^0 \left[\int_0^{t \wedge \tau_B} \mathbf{1}_{U_n}(X_s) \, ds \right], \end{aligned}$$

where

$$\bar{c}_2(\alpha, n) = |B_n| \cdot [2\delta_n - \delta'_n + 2y_n + 1 + (\delta_n + 1) \cdot \tan(A_n + B_n)]^{-\alpha-2}.$$

Letting $t \rightarrow \infty$, by the dominated convergence theorem, it follows

$$u_n(0) \geq \bar{c}_2(\alpha, n) \cdot \mathbb{E}^0 \left[\int_0^{\tau_B} \mathbf{1}_{U_n}(X_s) \, ds \right].$$

Using the estimate of the Green function from [2] there is $c(\alpha)$ so that

$$\begin{aligned}
\mathbb{E}^0 \left[\int_0^{\tau_B} \mathbf{1}_{U_n}(X_s) \, ds \right] &= \mathbb{E}^0 \left[\int_0^\infty \mathbf{1}_{\{X_s \in U_n, \tau_B > s\}} \, ds \right] \\
&= \int_0^\infty \mathbb{P}^0(X_s \in U_n, \tau_B > s) \, ds \\
&= \int_{U_n} G_B(0, y) \, dy \\
&\geq \int_{U_n} s(y) |y|^{\alpha-2} \, dy \\
&\geq c(\alpha) \cdot (1 - 2y_n)^\alpha \int_{U_n} |y|^{\alpha-2} \, dy \\
&\geq c(\alpha) \cdot (1 - 2y_n)^\alpha \int_{B(0, y_n/2)} |y|^{\alpha-2} \, dy \\
&\geq c(\alpha) \cdot (1 - 2y_n)^\alpha (y_n)^\alpha.
\end{aligned}$$

From here we obtain the lower bound

$$(2.24) \quad u_n(0) \geq c_2(\alpha, n) \cdot (1 - 2y_n)^\alpha (y_n)^\alpha,$$

where

$$c_2(\alpha, n) = c(\alpha) |B_n| \cdot [2\delta_n - \delta'_n + 2y_n + 1 + (\delta_n + 1) \cdot \tan(A_n + B_n)]^{-\alpha-2}.$$

Combining inequalities (2.23) and (2.24), we obtain:

$$(2.25) \quad \frac{u_n(w_0)}{u_n(0)} \leq c(\alpha) \cdot a_n b_n,$$

where

$$\begin{aligned}
a_n &= \left(\frac{2\delta_n - \delta'_n + 2y_n + 1}{(\delta_n + 1) \cdot \tan(A_n + B_n)} + 1 \right)^{\alpha+2}, \\
b_n &= \frac{\delta_n + x_n}{y_n^\alpha \cdot (1 - 2y_n)^\alpha}.
\end{aligned}$$

Due to the construction,

$$(2.26) \quad \lim_{n \rightarrow \infty} a_n = 1.$$

Using (2.3) and (2.4), we obtain

$$(2.27) \quad \limsup_{n \rightarrow \infty} \frac{\delta_n + x_n}{y_n} < \infty.$$

Since the proof of (2.27) is similar to the proof of Lemma 4.7 (Appendix), we skip it.

Since $0 < \alpha < 1$,

$$\begin{aligned}
b_n &= \frac{\delta_n + x_n}{y_n^\alpha \cdot (1 - 2y_n)^\alpha} \\
&= \frac{1}{(1 - 2y_n)^\alpha} \cdot \frac{\delta_n + x_n}{y_n} \cdot y_n^{1-\alpha}.
\end{aligned}$$

Therefore,

$$(2.28) \quad \lim_{n \rightarrow \infty} b_n = 0.$$

To conclude, (2.25), (2.26) and (2.28) imply

$$\frac{u_n(0)}{u_n(w_0)}$$

can be made as large as we like by taking n large enough. This implies that the Harnack inequality for X is not possible. $\square$

Remark 2.15. Notice that for $1 < \alpha < 2$, we have

$$\frac{u_n(0)}{u_n(w_0)} \geq c(\alpha) \cdot \frac{1}{a_n} \cdot \frac{1}{b_n},$$

where

$$\frac{1}{b_n} = (1 - 2y_n)^\alpha \cdot \frac{y_n}{\delta_n + x_n} \cdot y_n^{\alpha-1}.$$

Notice that from here, we have

$$\lim_{n \rightarrow \infty} \frac{1}{b_n} = 0,$$

therefore the proof breaks down.

Remark 2.16. In connection with the article [2], there it was shown that as in our case ($d = 2$), for $1 < \alpha < 2$ Harnack inequality holds (see Corollary 13 of the aforementioned article).

3. WEAK HARNACK INEQUALITY

Definition 3.1. *The weak Harnack inequality* for a symmetric α -stable Lévy processes X holds if there is a constant $C \geq 1$ such that for every non-negative function $u: \mathbb{R}^d \rightarrow \mathbb{R}$, which is harmonic in B_1 with respect to X , the inequality

$$\|u\|_{L^1(B_{1/2})} \leq C \cdot \inf_{B_{1/4}} u$$

holds.

We continue with the definitions regarding the spherical part of the Lévy measure (see [3] (cf. [8])).

Definition 3.2. A measure λ on $\mathbb{R}^d$ is called *degenerate* if there is a proper linear subspace M of $\mathbb{R}^d$ such that $Spt(\lambda) \subset M$, where $Spt(\lambda)$ denotes the support of the measure μ .

A measure λ is called *non-degenerate* if it is not degenerate.

Definition 3.3. A measure μ on $\mathbb{S}^{d-1}$ is called a *spectral measure* if it is positive, finite, non-degenerate and symmetric.

For the equivalence of the non-degeneracy of the measure μ and the condition

$$(3.1) \quad \Phi(u) \geq c \cdot |u|^\alpha, \quad u \in \mathbb{R}^d,$$

where $c = c(\alpha)$ is a positive constant and Φ is as in (1.2), see [8].

Theorem 3.4 (Weak Harnack Inequality). *Let X be a symmetric α -stable Lévy process in $\mathbb{R}^d$, $d \geq 2$, with index of stability $\alpha \in (0, 2)$ and the characteristic function of the form*

$$\mathbb{E}^0 e^{iu \cdot X_t} = e^{-t\Phi(u)}, \quad u \in \mathbb{R}^d, \quad t \geq 0,$$

where the characteristic exponent is given by

$$\Phi(u) = \int_{\mathbb{S}^{d-1}} |u \cdot \xi|^\alpha \mu(d\xi),$$

and μ is a spectral measure. Furthermore, let μ be absolutely continuous with respect to the uniform measure σ on the sphere $\mathbb{S}^{d-1}$ and denote by f_μ its density. Assume that there is a positive constant m such that

$$0 \leq f_\mu(\xi) \leq m, \quad \xi \in \mathbb{S}^{d-1}.$$

Then the weak Harnack inequality for X holds.

Lemma 3.5. *Let $\alpha \in (0, 2)$, $d \geq 2$. There is a constant $c_1 = c_1(\alpha, d)$ such that for every $|w| < 3/4$, the inequality*

$$\int_{B_{1/2}} G_{B_{3/4}}(x, w) \, dx \leq c_1$$

holds.

Proof. By the inequality $p_D \leq p$ and the estimate of the transition density p for small times (see e.g. [10], Theorem 1), it follows:

$$\begin{aligned}
& \int_{B_{1/2}} G_{B_{3/4}}(x, w) \, dx \\
&= \int_{B_{1/2}} \left[\int_0^\infty p_{B_{3/4}}(t, x, w) \, dt \right] \, dx \\
&\leq \int_{B_{1/2}} \left[\int_0^\infty p(t, x, w) \, dt \right] \, dx \\
&\leq c(\alpha, d) \int_{B_{1/2}} \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|x-w|^{\alpha+d}} \right) \, dt \right] \, dx \\
&\quad + \int_{B_{1/2}} \left[\int_1^\infty t^{-d/\alpha} \cdot p(1, t^{-1/\alpha}x, t^{-1/\alpha}w) \, dt \right] \, dx \\
&\leq c(\alpha, d) \int_{B_{1/2}} \left[\int_0^{|x-w|^\alpha \wedge 1} \frac{t}{|x-w|^{\alpha+d}} \, dt \right] \, dx \\
&\quad + c(\alpha, d) \int_{B_{1/2}} \left[\int_{|x-w|^\alpha \wedge 1}^1 t^{-d/\alpha} \, dt \right] \, dx \\
&\quad + c(\alpha, d) \left[\int_{B_{1/2}} \int_1^\infty t^{-d/\alpha} \, dt \right] \, dx \\
&\leq c(\alpha, d) \left[\int_{B_{1/2}} |x-w|^{\alpha-d} \, dx \right. \\
&\quad \left. + \frac{\alpha}{d-\alpha} \int_{B_{1/2}} \left[(|x-w|^\alpha \wedge 1)^{\frac{\alpha-d}{\alpha}} - 1 \right] \, dx \right] \\
&\quad + c(\alpha, d) \frac{\alpha}{d-\alpha} |B_{1/2}| \\
&\leq c(\alpha, d).
\end{aligned}$$

□

Lemma 3.6. *Let $\alpha \in (0, 2), d \geq 2$. There exist $\delta_1 = \delta_1(\alpha, d) > 0$ and $c_2 = c_2(\alpha, d, \delta_1)$ such that for every $|\bar{x}| < 1/4$ and every $w \in B(\bar{x}, \delta_1)$ the inequality*

$$G_{B_{3/4}}(\bar{x}, w) \geq c_2$$

holds.

Proof. Using

$$(3.2) \quad G_{B_{3/4}}(\bar{x}, w) = G(\bar{x}, w) - \mathbb{E}^{\bar{x}}[G(X_{\tau_{B_{3/4}}}, w)],$$

in order to prove the lemma, we compute the estimates for $G(\bar{x}, w)$ from below and $\mathbb{E}^{\bar{x}}[G(X_{\tau_{B_{3/4}}}, w)]$ from above. Using the heat kernel estimates for small times ([10],

Theorem 1), we obtain

$$\begin{aligned}
& \mathbb{E}^{\bar{x}}[G(X_{\tau_{B_{3/4}}}, w)] \\
&= \int_{B_{3/4}^c} G(u, w) P_{B_{3/4}}(\bar{x}, u) \, du \\
&= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_0^\infty p(t, u, w) \, dt \right] \, du \\
&= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_0^1 p(t, u, w) \, dt \right] \, du \\
&\quad + \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_1^\infty p(t, u, w) \, dt \right] \, du \\
&\leq c(\alpha, d) \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) \, dt \right] \, du \\
&\quad + \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_1^\infty t^{-d/\alpha} \cdot p(1, t^{-1/\alpha}u, t^{-1/\alpha}w) \, dt \right] \, du \\
&\leq c(\alpha, d) \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) \, dt \right] \, du \\
&\quad + c(\alpha, d) \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_1^\infty t^{-d/\alpha} \, dt \right] \, du \\
&\leq c(\alpha, d) \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) \, dt \right] \, du \\
(3.3) \quad &+ c(\alpha, d).
\end{aligned}$$

Examining the integral in (3.3) more closely, we obtain:

$$\begin{aligned}
& \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) \, dt \right] \, du \\
&= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|<1\}}(u) \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) \, dt \right] \, du \\
&\quad + \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|>1\}}(u) \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) \, dt \right] \, du \\
(3.4) \quad &= I_1 + I_2.
\end{aligned}$$

$$\begin{aligned}
I_1 &= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|<1\}}(u) \\
&\quad \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) dt \right] du \\
&= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|<1\}}(u) |w-u|^{-\alpha-d} \left[\int_0^{|w-u|^\alpha} t dt \right] du \\
&\quad + \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|<1\}}(u) \left[\int_{|w-u|^\alpha}^1 t^{-d/\alpha} dt \right] du \\
&= c_1 \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|<1\}}(u) |w-u|^{\alpha-d} du \\
&\quad + c_2 \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|<1\}}(u) \frac{\alpha}{d-\alpha} (|w-u|^{\alpha-d} - 1) du \\
(3.5) \quad &\leq c_1(\alpha, d, \bar{\delta}_1),
\end{aligned}$$

where in the last inequality we have used $w \in B(\bar{x}, \bar{\delta}_1)$, for $\bar{\delta}_1 > 0$ small enough.

$$\begin{aligned}
I_2 &= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|>1\}}(u) \\
&\quad \left[\int_0^1 \left(t^{-d/\alpha} \wedge \frac{t}{|u-w|^{\alpha+d}} \right) dt \right] du \\
&= \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|>1\}}(u) |w-u|^{-\alpha-d} \left[\int_0^1 t dt \right] du \\
&= c \int_{B_{3/4}^c} P_{B_{3/4}}(\bar{x}, u) \mathbf{1}_{\{|w-u|>1\}}(u) |w-u|^{-\alpha-d} du \\
(3.6) \quad &\leq c_2.
\end{aligned}$$

In conclusion, by (3.3), (3.4), (3.5) and (3.6), we obtain

$$(3.7) \quad \mathbb{E}^{\bar{x}}[G(X_{\tau_{B_{3/4}}}, w)] \leq \bar{c}_1(\alpha, d, \bar{\delta}_1),$$

for all $w \in B(\bar{x}, \bar{\delta}_1)$ and $\bar{\delta}_1 > 0$ small enough.

To estimate $G(\bar{x}, w)$ from below, we use the continuity of the potential density (see [11]).

Due to

$$p(1, 0) \geq c > 0,$$

by continuity of $p(1, \cdot)$ in $x = 0$, there is $R > 0$ such that $p(1, x) > \frac{1}{2} \cdot p(1, 0)$, for all $|x| < R$.

Furthermore, for $|\xi| = 1$, since:

$$\begin{aligned} G(0, \xi) &\geq \int_{R^{-\alpha}}^{\infty} p(t, \xi) \, dt \\ &= \int_{R^{-\alpha}}^{\infty} t^{-d/\alpha} \cdot p\left(1, \frac{\xi}{t^{1/\alpha}}\right) \, dt > \frac{1}{2} \cdot \int_{R^{-\alpha}}^{\infty} t^{-d/\alpha} \cdot p(1, 0) \, dt \\ &= c_1(\alpha, d) > 0, \end{aligned}$$

we obtain

$$(3.8) \quad G(0, \xi) \geq c_1(\alpha, d).$$

For $|x| \neq 0$, by scaling and (3.8)

$$G(0, x) = |x|^{\alpha-d} \cdot G\left(0, \frac{x}{|x|}\right) \geq c_1 \cdot |x|^{\alpha-d}.$$

Therefore,

$$(3.9) \quad G(\bar{x}, w) \geq \bar{c}_2 \cdot |\bar{x} - w|^{\alpha-d}.$$

Now, choose δ_1 such that $\delta_1 < (\bar{c}_2/(\bar{c}_1 + c))^{\frac{1}{\alpha-d}} \wedge \bar{\delta}_1$, where $c > 0$. Then for every $|\bar{x}| < 1/4$ and $w \in B(\bar{x}, \delta_1)$, combining (3.2), (3.7) and (3.9), we obtain

$$G_{B_{3/4}}(\bar{x}, w) \geq \bar{c}_2 \cdot |\bar{x} - w|^{\alpha-d} - \bar{c}_1 \geq c > 0.$$

Define $c_2 = c$ and now the statement follows. $\square$

Remark 3.7. Notice that, according to the Lemma 3.5 and Lemma 3.6, there are $\tilde{c} = \tilde{c}(\alpha, d)$ and $\delta_1 = \delta_1(\alpha, d)$ such that for every $\bar{x} \in B_{1/4}$ and for every $w \in B(\bar{x}, \delta_1)$ the inequality

$$\int_{B_{1/2}} G_{B_{3/4}}(x, w) \, dx \leq \tilde{c} \cdot G_{B_{3/4}}(\bar{x}, w)$$

holds.

Lemma 3.8. *For $\alpha \in (0, 2)$ and $d \geq 2$, let $\delta_1 > 0$ be as in Lemma 3.6. There is a constant $c_3 = c_3(\alpha, d)$ such that for every $\bar{x} \in B_{1/4}$ and every $\tilde{u} \in B(0, 3/4) \setminus B(\bar{x}, \delta_1)$ the inequality*

$$\int_{B_{1/2}} G_{B_{3/4}}(x, \tilde{u}) \, dx \leq c_3 \cdot G_{B_{3/4}}(\bar{x}, \tilde{u})$$

holds.

Proof. The proof relies on the maximum principle (cf. [7]). We use the fact that $G_D(\bar{x}, \cdot)$ is regular harmonic in $D \setminus B(\bar{x}, \varepsilon)$ with respect to X for every $\varepsilon > 0$ (cf. [1]).

$$\begin{aligned}
\int_{B_{1/2}} G_{B_{3/4}}(x, \tilde{u}) \, dx &= \int_{B_{1/2}} \mathbb{E}^{\tilde{u}}[G_{B_{3/4}}(x, X_{\tau_{B_{3/4} \setminus B(\bar{x}, \delta_1)}})] \, dx \\
&= \int_{B_{1/2}} \left[\int_{\mathbb{R}^2 \setminus (B_{3/4} \setminus B(\bar{x}, \delta_1))} G_{B_{3/4}}(x, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dz \right] dx \\
&= \int_{\mathbb{R}^2 \setminus (B_{3/4} \setminus B(\bar{x}, \delta_1))} \left[\int_{B_{1/2}} G_{B_{3/4}}(x, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dx \right] dz \\
&= \int_{B(\bar{x}, \delta_1)} \left[\int_{B_{1/2}} G_{B_{3/4}}(x, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dx \right] dz \\
&\quad + \int_{\mathbb{R}^2 \setminus B_{3/4}} \left[\int_{B_{1/2}} G_{B_{3/4}}(x, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dx \right] dz \\
&= \int_{B(\bar{x}, \delta_1)} \left[\int_{B_{1/2}} G_{B_{3/4}}(x, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dx \right] dz \\
&\stackrel{Rem.3.7}{\leq} \tilde{c} \int_{B(\bar{x}, \delta_1)} G_{B_{3/4}}(\bar{x}, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dz \\
&= \tilde{c} \int_{\mathbb{R}^2 \setminus (B_{3/4} \setminus B(\bar{x}, \delta_1))} G_{B_{3/4}}(\bar{x}, z) P_{B_{3/4} \setminus B(\bar{x}, \delta_1)}(\tilde{u}, z) \, dz \\
&= \tilde{c} \cdot \mathbb{E}^{\tilde{u}}[G_{B_{3/4}}(\bar{x}, X_{\tau_{B_{3/4} \setminus B(\bar{x}, \delta_1)}})] \\
&= \tilde{c} \cdot G_{B_{3/4}}(\bar{x}, \tilde{u}).
\end{aligned}$$

Define $c_3 = \tilde{c}$ and the lemma follows. $\square$

Proof of Theorem 3.4.

$$\begin{aligned}
& \|u\|_{L^1(B_{1/2})} \\
&= \int_{B_{1/2}} u(x) \, dx \\
&= \int_{B_{1/2}} \mathbb{E}^x[u(X_{\tau_{B_{3/4}}})] \, dx \\
&= \int_{B_{1/2}} \left[\int_{(\bar{B}_{3/4})^c} u(y) P_{B_{3/4}}(x, y) \, dy \right] dx \\
&= \int_{B_{1/2}} \left[\int_{(\bar{B}_{3/4})^c} u(y) \left[\int_{B(y, 3/4)} f_\nu(z) G_{B_{3/4}}(x, y - z) \, dz \right] dy \right] dx \\
(3.10) \quad &= \int_{(\bar{B}_{3/4})^c} u(y) \left[\int_{B(y, 3/4)} f_\nu(z) \left[\int_{B_{1/2}} G_{B_{3/4}}(x, y - z) \, dx \right] dz \right] dy
\end{aligned}$$

By Remark 3.7 and Lemma 3.8 there is a constant $c = c(\alpha, d)$ such that for every $\bar{x} \in B_{1/4}$

$$\begin{aligned}
(3.10) \quad &\leq c \cdot \int_{(\bar{B}_{3/4})^c} u(y) \left[\int_{B(y, 3/4)} \mathbf{1}_{B(\bar{x}, \delta_1)}(y - z) \right. \\
&\quad \left. f_\nu(z) \cdot G_{B_{3/4}}(\bar{x}, y - z) \, dz \right] dy \\
&+ c \cdot \int_{(\bar{B}_{3/4})^c} u(y) \left[\int_{B(y, 3/4)} \mathbf{1}_{B(\bar{x}, \delta_1)^c}(y - z) \right. \\
&\quad \left. f_\nu(z) \cdot G_{B_{3/4}}(\bar{x}, y - z) \, dz \right] dy \\
&= c \cdot \int_{(\bar{B}_{3/4})^c} u(y) \left[\int_{B(y, 3/4)} f_\nu(z) G_{B_{3/4}}(\bar{x}, y - z) \, dz \right] dy \\
&= c \cdot \int_{(\bar{B}_{3/4})^c} u(y) P_{B_{3/4}}(\bar{x}, y) \, dy \\
&= c \cdot \mathbb{E}^{\bar{x}}[u(X_{\tau_{B_{3/4}}})] \\
&= c \cdot u(\bar{x}).
\end{aligned}$$

Since the inequality

$$\|u\|_{L^1(B_{1/2})} \leq c \cdot u(\bar{x}),$$

holds for any $\bar{x} \in B_{1/4}$, the proof is finished. $\square$

4. APPENDIX

Lemma 4.1. *Let:*

- (1) $\alpha_k = 2^{-k}$, $\beta_k = \frac{1}{k(k+1)}$,
- (2) $\alpha_k = a^{-k}$, $\beta_k = b^{-k}$, $1 < b \leq a$,
- (3) $\alpha_k = \frac{1}{k(k+1)}$, $\beta_k = k^{-1-\delta}$, $\delta > 0$,

$$(4) \quad \alpha_k = 2^{-(k^3)}, \quad \beta_k = 2^{-(k^2)} \\ \alpha_k = a^{-(k^3)}, \quad \beta_k = b^{-(k^2)}, \quad a > 1, b > 1.$$

Then none of these sequences does not satisfy the property (2.3) from the Lemma 2.3.

Proof. Notice that

$$\lim_{n \rightarrow \infty} \frac{\tan(A_n + B_{n+1})}{\tan(A_n + B_n)} = \infty$$

implies

$$(4.1) \quad \lim_{n \rightarrow \infty} \frac{\cos(A_n + B_{n+1})}{\cos(A_n + B_n)} = 0.$$

(1) Set

$$\alpha_k = 2^{-k}, \quad \beta_k = \frac{1}{k(k+1)}.$$

Since $A_n = 1 - 2^{-n}$, $B_n = \frac{n}{n+1}$ and $B_{n+1} = B_n + \beta_{n+1}$, we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\cos(A_n + B_{n+1})}{\cos(A_n + B_n)} &= \lim_{n \rightarrow \infty} \frac{\frac{d}{dn}(A_n + B_{n+1})}{\frac{d}{dn}(A_n + B_n)} \\ &= 1 + \lim_{n \rightarrow \infty} \frac{\frac{d}{dn}\beta_{n+1}}{\frac{d}{dn}(A_n + B_n)} \\ &= 1 + \lim_{n \rightarrow \infty} \frac{\frac{-2n-3}{(n+1)^2(n+2)^2}}{\ln 2 \cdot 2^{-n} + (n+1)^{-2}} \\ &= 1 + 0 = 1. \end{aligned}$$

This means that the condition (4.1) is not fulfilled.

(2) Set

$$\alpha_k = a^{-k}, \quad \beta_k = b^{-k}, \quad 1 < b \leq a.$$

Since $A_n = \frac{1-a^{-n}}{a-1}$, $B_n = \frac{1-b^{-n}}{b-1}$, we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\cos(A_n + B_{n+1})}{\cos(A_n + B_n)} &= \lim_{n \rightarrow \infty} \frac{\frac{d}{dn}(A_n + B_{n+1})}{\frac{d}{dn}(A_n + B_n)} \\ &= 1 + \lim_{n \rightarrow \infty} \frac{\frac{d}{dn}\beta_{n+1}}{\frac{d}{dn}(A_n + B_n)} \\ &= 1 + \lim_{n \rightarrow \infty} \frac{-\ln b \cdot b^{-(n+1)}}{\frac{\ln a}{a-1} a^{-n} + \frac{\ln b}{b-1} b^{-n}} \\ &= 1 + \lim_{n \rightarrow \infty} \frac{-\ln b \cdot b^{-1}}{\frac{\ln a}{a-1} \left(\frac{b}{a}\right)^n + \frac{\ln b}{b-1}}. \end{aligned}$$

If $a = b$, the last expression is equal to

$$1 - \frac{\ln a \cdot a^{-1}}{2 \frac{\ln a}{a-1}} = 1 - \frac{a-1}{2a} = \frac{a+1}{2a}.$$

If $b < a$, we obtain

$$1 - \frac{\ln b \cdot b^{-1}}{\frac{\ln b}{b-1}} = 1 - \frac{b-1}{b} = \frac{1}{b}.$$

In conclusion, the condition (4.1) is not fulfilled.

Remark 4.2. In the examples that follow, we will proceed in the following way.

We use the equality

$$\frac{\cos(A_n + B_{n+1})}{\cos(A_n + B_n)} = \frac{\sin(\frac{\pi}{2} - A_n - B_{n+1})}{\sin(\frac{\pi}{2} - A_n - B_n)},$$

and the estimates

$$(4.2) \quad c_1 \cdot \frac{\frac{\pi}{2} - A_n - B_{n+1}}{\frac{\pi}{2} - A_n - B_n} \leq \frac{\sin(\frac{\pi}{2} - A_n - B_{n+1})}{\sin(\frac{\pi}{2} - A_n - B_n)}$$

$$(4.3) \quad \frac{\sin(\frac{\pi}{2} - A_n - B_{n+1})}{\sin(\frac{\pi}{2} - A_n - B_n)} \leq c_2 \cdot \frac{\frac{\pi}{2} - A_n - B_{n+1}}{\frac{\pi}{2} - A_n - B_n},$$

for some constants $c_1, c_2 > 0$.

Examining the lower (or upper) bound in the expression (4.2) (or (4.3)), we see

$$(4.4) \quad \begin{aligned} \frac{\frac{\pi}{2} - A_n - B_n - \beta_{n+1}}{\frac{\pi}{2} - A_n - B_n} &= 1 - \frac{\beta_{n+1}}{\frac{\pi}{2} - A_n - B_n} \\ &= 1 - \frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k}. \end{aligned}$$

Now, we can use integral test for convergence to estimate series

$$\sum_{k=n+1}^{\infty} \alpha_k$$

and

$$\sum_{k=n+1}^{\infty} \beta_k.$$

(3) Let

$$\alpha_k = \frac{1}{k(k+1)}, \quad \beta_k = k^{-1-\delta},$$

for some $\delta > 0$. In this example we will use integral test.

Notice,

$$\sum_{k=n+1}^{\infty} \alpha_k = \frac{1}{n+1}.$$

The integral test implies

$$\int_{n+1}^{\infty} \beta(x) dx \leq \sum_{k=n+1}^{\infty} \beta_k \leq \beta_{n+1} + \int_{n+1}^{\infty} \beta(x) dx,$$

which is equivalent to

$$\begin{aligned} \sum_{k=n+1}^{\infty} \alpha_k + \int_{n+1}^{\infty} \beta(x) dx &\leq \sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k, \\ \sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k &\leq \sum_{k=n+1}^{\infty} \alpha_k + \beta_{n+1} + \int_{n+1}^{\infty} \beta(x) dx. \end{aligned}$$

These inequalities yield

$$\begin{aligned} \frac{1}{\sum_{k=n+1}^{\infty} \alpha_k + \beta_{n+1} + \int_{n+1}^{\infty} \beta(x) dx} &\leq \frac{1}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k}, \\ \frac{1}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k} &\leq \frac{1}{\sum_{k=n+1}^{\infty} \alpha_k + \int_{n+1}^{\infty} \beta(x) dx}. \end{aligned}$$

Multiplying the two inequalities by β_{n+1} , we obtain the lower and the upper bound for the expression in (4.4),

$$\frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \beta_{n+1} + \int_{n+1}^{\infty} \beta(x) dx} \leq \frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k}$$

and

$$\frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k} \leq \frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \int_{n+1}^{\infty} \beta(x) dx}.$$

Since

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=n+1}^{\infty} \alpha_k}{\beta_{n+1}} = \infty,$$

it follows

$$\lim_{n \rightarrow \infty} \frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \int_{n+1}^{\infty} \beta(x) dx} = 0$$

and consequently

$$\lim_{n \rightarrow \infty} \frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k} = 0.$$

This means, by (4.4),

$$\lim_{n \rightarrow \infty} \frac{\frac{\pi}{2} - A_n - B_{n+1}}{\frac{\pi}{2} - A_n - B_n} = 1 - 0 = 1,$$

which implies the condition (4.1) is not fulfilled.

(4) Set

$$\alpha_k = 2^{-(k^3)}, \quad \beta_k = 2^{-(k^2)}.$$

As in the second example, we use the integral test. Similarly, it implies

$$\begin{aligned} \int_{n+1}^{\infty} \alpha(x) dx + \int_{n+1}^{\infty} \beta(x) dx &\leq \sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k, \\ \sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k &\leq \alpha_{n+1} + \int_{n+1}^{\infty} \alpha(x) dx + \beta_{n+1} + \int_{n+1}^{\infty} \beta(x) dx, \end{aligned}$$

which yields

$$\begin{aligned} &\frac{1}{\alpha_{n+1} + \int_{n+1}^{\infty} \alpha(x) dx + \beta_{n+1} + \int_{n+1}^{\infty} \beta(x) dx} \\ &\leq \frac{1}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k}, \\ (4.5) \quad &\frac{1}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k} \leq \frac{1}{\int_{n+1}^{\infty} \alpha(x) dx + \int_{n+1}^{\infty} \beta(x) dx}. \end{aligned}$$

We will only observe the inequality (4.5). Multiplying (4.5) by β_{n+1} , it follows

$$\frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k} \leq \frac{\beta_{n+1}}{\int_{n+1}^{\infty} \alpha(x) dx + \int_{n+1}^{\infty} \beta(x) dx}.$$

Since

$$\frac{\beta_{n+1}}{\int_{n+1}^{\infty} \alpha(x) dx + \int_{n+1}^{\infty} \beta(x) dx} \leq \frac{\beta_{n+1}}{\int_{n+1}^{\infty} \beta(x) dx},$$

it is enough to show

$$\lim_{n \rightarrow \infty} \frac{\beta_{n+1}}{\int_{n+1}^{\infty} \beta(x) dx} = 0.$$

Notice that then

$$\lim_{n \rightarrow \infty} \frac{\beta_{n+1}}{\sum_{k=n+1}^{\infty} \alpha_k + \sum_{k=n+1}^{\infty} \beta_k} = 0,$$

which implies the condition (4.1) is not fulfilled.

Using polar coordinates we obtain

$$\begin{aligned} \int_{n+1}^{\infty} \beta(x) dx &= \int_{n+1}^{\infty} 2^{-(x^2)} dx \\ &= \frac{1}{2} \sqrt{\frac{\pi}{\ln 2}} \cdot 2^{-\frac{1}{2}(n+1)^2}, \end{aligned}$$

which implies

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\beta_{n+1}}{\int_{n+1}^{\infty} \beta(x) dx} &= \lim_{n \rightarrow \infty} \frac{2^{-(n+1)^2}}{\frac{1}{2} \sqrt{\frac{\pi}{\ln 2}} \cdot 2^{-\frac{1}{2}(n+1)^2}} \\ &= \lim_{n \rightarrow \infty} \frac{2^{-\frac{1}{2}(n+1)^2}}{\frac{1}{2} \sqrt{\frac{\pi}{\ln 2}}} \\ &= 0, \end{aligned}$$

and that is what we wanted to prove. $\square$

Remark 4.3. Notice that only in the very end the explicit formula for β_n has been used. This means that if

$$\beta_k = b^{-(k^2)}, \quad b > 1,$$

the same reasoning applies.

In this case, we would have

$$\begin{aligned} \int_{n+1}^{\infty} \beta(x) dx &= \int_{n+1}^{\infty} b^{-(x^2)} dx \\ &= \frac{1}{2} \sqrt{\frac{\pi}{\ln b}} \cdot b^{-\frac{1}{2}(n+1)^2} \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\beta_{n+1}}{\int_{n+1}^{\infty} \beta(x) dx} &= \lim_{n \rightarrow \infty} \frac{b^{-(n+1)^2}}{\frac{1}{2} \sqrt{\frac{\pi}{\ln b}} \cdot b^{-\frac{1}{2}(n+1)^2}} \\ &= \lim_{n \rightarrow \infty} \frac{b^{-\frac{1}{2}(n+1)^2}}{\frac{1}{2} \sqrt{\frac{\pi}{\ln b}}} \\ &= 0. \end{aligned}$$

Lemma 4.4. *Let the quantities δ_n , I_n , P_n , P'_n , δ_n , x_n , y_n , c_n and d_n be as in the Proposition 2.2. Then:*

- (1) $d_n = (\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1$;
- (2) *all quantities above are positive and $c_n < d_n$.*

Proof. (1) By the definition (2.10),

$$\delta_n = \frac{I_n(1 + \tan(A_n + B_n)) \cdot (1 + \tan(A_{n+1} + B_{n+1}))}{P_n^2(1 + \tan(A_n + B_{n+1})) + P'_n I_n(1 + \tan(A_{n+1} + B_{n+1}))}.$$

This expression is equivalent to

$$\begin{aligned} &I_n(1 + \tan(A_n + B_n)) \cdot (1 + \tan(A_{n+1} + B_{n+1})) \\ &= \delta_n P'_n I_n \cdot (1 + \tan(A_{n+1} + B_{n+1})) + \delta_n P_n^2 \cdot (1 + \tan(A_n + B_{n+1})), \end{aligned}$$

from where

$$\begin{aligned} &I_n(1 + \tan(A_n + B_n)) \cdot (1 + \tan(A_{n+1} + B_{n+1})) \\ &= \delta_n P'_n I_n \cdot (1 + \tan(A_{n+1} + B_{n+1})) \\ &\quad + \delta_n P_n(1 + \tan(A_n + B_{n+1})) \cdot (I_n - 2(1 + \tan(A_n + B_{n+1}))) \end{aligned}$$

follows. Therefore,

$$\begin{aligned} & 2\delta_n P_n \cdot (1 + \tan(A_n + B_{n+1}))^2 \\ &= \delta_n P_n I_n \cdot (1 + \tan(A_n + B_{n+1})) \\ &+ I_n (1 + \tan(A_{n+1} + B_{n+1})) \cdot (\delta_n P'_n - \tan(A_n + B_n) - 1), \end{aligned}$$

and eventually

$$\begin{aligned} & 2\delta_n \frac{P_n}{I_n} \cdot (1 + \tan(A_n + B_{n+1})) = \\ &= \delta_n P_n + \frac{1 + \tan(A_{n+1} + B_{n+1})}{1 + \tan(A_n + B_{n+1})} \cdot (\delta_n P'_n - \tan(A_n + B_n) - 1). \end{aligned}$$

By the definition of x_n (2.11), the last equality yields

$$\begin{aligned} & x_n \cdot \tan(A_n + B_{n+1}) - 1 = \\ &= \delta_n P_n + \frac{1 + \tan(A_{n+1} + B_{n+1})}{1 + \tan(A_n + B_{n+1})} \cdot (\delta_n P'_n - \tan(A_n + B_n) - 1). \end{aligned}$$

Therefore,

$$\frac{1 + \tan(A_n + B_n) - \delta_n P'_n}{1 + \tan(A_n + B_{n+1})} = \frac{\delta_n P_n - x_n \cdot \tan(A_n + B_{n+1}) + 1}{1 + \tan(A_{n+1} + B_{n+1})}.$$

Notice that by (2.12) the left-hand side is y_n , hence

$$y_n = \frac{\delta_n P_n - x_n \cdot \tan(A_n + B_{n+1}) + 1}{1 + \tan(A_{n+1} + B_{n+1})}.$$

From here, by the definition of P_n (2.8),

$$\begin{aligned} & (\delta_n - y_n) \cdot \tan(A_{n+1} + B_{n+1}) - y_n \\ &= (\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1. \end{aligned}$$

Consequently, by the definition of d_n (2.14), the last equality implies

$$d_n = (\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1,$$

and hence the first part of the Lemma.

- (2) Using the definition of δ_n (2.10) one sees that $y_n > 0$. Namely, the nominator of y_n is positive if and only if

$$1 + \tan(A_n + B_n) - \delta_n \cdot P'_n > 0,$$

which is equivalent to

$$\frac{1 + \tan(A_n + B_n)}{P'_n} > \delta_n.$$

By the definition of δ_n , the condition above is fulfilled if and only if

$$\begin{aligned} & \frac{1 + \tan(A_n + B_n)}{P'_n} \\ & > \frac{I_n \cdot (1 + \tan(A_n + B_n)) \cdot (1 + \tan(A_{n+1} + B_{n+1}))}{H_n}, \end{aligned}$$

which is equivalent to

$$H_n > P'_n \cdot I_n \cdot (1 + \tan(A_{n+1} + B_{n+1})).$$

Using (2.15), we see that it holds.

Consequently, c_n is also positive.

Next we show that $c_n < d_n$, which also implies that $d_n > 0$. Since

$$I_n > 0,$$

by the definition of P_n (2.8), the inequality

$$2 \cdot (1 + \tan(A_{n+1} + B_{n+1})) > P_n$$

holds.

Therefore,

$$\begin{aligned} & 2 \cdot P_n(1 + \tan(A_n + B_{n+1})) \cdot (1 + \tan(A_{n+1} + B_{n+1})) \\ & > P_n^2(1 + \tan(A_n + B_{n+1})). \end{aligned}$$

Using the expression for H_n (2.15),

$$\begin{aligned} & 2 \cdot P_n(1 + \tan(A_n + B_{n+1})) \cdot (1 + \tan(A_{n+1} + B_{n+1})) \\ & + P'_n I_n \cdot (1 + \tan(A_{n+1} + B_{n+1})) > H_n, \end{aligned}$$

which, by the definition of δ_n (2.10) yields

$$\delta_n > \frac{I_n \cdot (1 + \tan(A_n + B_n))}{2 \cdot P_n(1 + \tan(A_n + B_{n+1})) + P'_n I_n}.$$

Therefore,

$$2\delta_n \cdot P_n(1 + \tan(A_n + B_{n+1})) > (1 + \tan(A_n + B_n) - \delta_n P'_n) \cdot I_n,$$

which implies

$$2\delta_n \cdot \frac{1 + \tan(A_n + B_{n+1})}{\tan(A_n + B_{n+1})} \cdot \frac{P_n}{I_n} > \frac{1 + \tan(A_n + B_n) - \delta_n P'_n}{\tan(A_n + B_{n+1})}.$$

Consequently, by the definition of x_n (2.11)

$$x_n > \frac{2 + \tan(A_n + B_n) - \delta_n P'_n}{\tan(A_n + B_{n+1})}.$$

Hence, using the definition for y_n (2.12),

$$\frac{x_n \cdot \tan(A_n + B_{n+1}) - 1}{1 + \tan(A_n + B_{n+1})} > y_n.$$

Therefore,

$$(\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1 > (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n.$$

By the first part of the Lemma and the definition of c_n (2.13), the equality

$$d_n > c_n,$$

holds.

Hence $d_n > 0$ and the proof is finished. $\square$

Lemma 4.5. *Let $I_n, J_n, P_n, P'_n, \delta_n, x_n, y_n, c_n, d_n, U_n, \delta'_n, S_n = (x_{S_n}, y_{S_n})$ and r_n be as in the Proposition 2.3. Then:*

$$(1) \ B(S_n, r_n) \subset Q_n := (\delta'_n, \delta_n) \times (c_n, c_n + (\delta_n - \delta'_n));$$

(2) the inequalities

$$\begin{aligned} c_n &\geq (\varepsilon + 1) \cdot \tan(A_n + B_n) + 1, \\ c_n + (\delta_n - \delta'_n) &\leq (\varepsilon + x_n) \cdot \tan(A_n + B_{n+1}) - 1, \end{aligned}$$

hold true, for some $\varepsilon > 0$ such that $\delta'_n < \varepsilon < \delta_n$.

Proof. (1) Let $z = (z_1, z_2) \in B(S_n, r_n)$. Clearly,

$$\begin{aligned} |z_1 - x_{S_n}| &< r_n, \\ |z_2 - y_{S_n}| &< r_n. \end{aligned}$$

If we show:

$$(4.6) \quad x_{S_n} - r_n = \delta'_n$$

$$(4.7) \quad x_{S_n} + r_n = \delta_n$$

$$(4.8) \quad y_{S_n} - r_n = c_n$$

$$(4.9) \quad y_{S_n} + r_n = c_n + (\delta_n - \delta'_n),$$

then the proof of the first part of the Lemma is finished.

Let us first check (4.6). The definitions of x_{S_n} (2.18), r_n (2.20), I_n (2.7), P_n (2.8) and δ'_n (2.16) imply

$$\begin{aligned} x_{S_n} - r_n &= \frac{\delta_n + y_n}{2} \cdot \frac{I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\ &\quad - \frac{1}{2} \cdot \frac{\delta_n P_n - y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\ &= \frac{\delta_n(I_n - P_n) + 2y_n I_n}{2(1 + \tan(A_{n+1} + B_{n+1}))} \\ &= \frac{\delta_n(1 + \tan(A_n + B_{n+1})) + y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\ &= \delta'_n. \end{aligned}$$

Let us show (4.7). Similarly, the definitions of x_{S_n} (2.18), r_n (2.20), I_n (2.7), P_n (2.8) and δ_n (2.10) imply

$$\begin{aligned} x_{S_n} + r_n &= \frac{\delta_n + y_n}{2} \cdot \frac{I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\ &\quad + \frac{1}{2} \cdot \frac{\delta_n P_n - y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\ &= \frac{\delta_n(I_n + P_n)}{2(1 + \tan(A_{n+1} + B_{n+1}))} \\ &= \frac{2\delta_n(1 + \tan(A_{n+1} + B_{n+1}))}{2(1 + \tan(A_{n+1} + B_{n+1}))} \\ &= \delta_n. \end{aligned}$$

Now, we check (4.8). Likewise, the definitions of y_{S_n} (2.19), r_n (2.20), I_n (2.7), J_n (2.17), P_n (2.8) and c_n (2.13) yield

$$\begin{aligned}
y_{S_n} - r_n &= \frac{\delta_n + y_n}{2} \cdot \frac{J_n}{1 + \tan(A_{n+1} + B_{n+1})} \\
&\quad - \frac{1}{2} \cdot \frac{\delta_n P_n - y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\
&= \frac{\delta_n(J_n - P_n) + y_n(I_n + J_n)}{2(1 + \tan(A_{n+1} + B_{n+1}))} \\
&= \delta_n \tan(A_n + B_{n+1}) + y_n \tan(A_n + B_{n+1}) + y_n \\
&= (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n \\
&= c_n.
\end{aligned}$$

Lastly, we check (4.9). The definitions of y_{S_n} (2.19), r_n (2.20), I_n (2.7), J_n (2.17) and P_n (2.8) imply

$$\begin{aligned}
y_{S_n} + r_n &= \frac{\delta_n + y_n}{2} \cdot \frac{J_n}{1 + \tan(A_{n+1} + B_{n+1})} \\
&\quad + \frac{1}{2} \cdot \frac{\delta_n \cdot P_n - y_n \cdot I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\
&= \frac{\delta_n \tan(A_{n+1} + B_{n+1})(1 + \tan(A_n + B_{n+1}))}{1 + \tan(A_{n+1} + B_{n+1})} \\
&\quad + \frac{y_n(\tan(A_n + B_{n+1}) \tan(A_{n+1} + B_{n+1}) - 1)}{1 + \tan(A_{n+1} + B_{n+1})} \\
&= \frac{\delta_n \tan(A_{n+1} + B_{n+1})(1 + \tan(A_n + B_{n+1}))}{1 + \tan(A_{n+1} + B_{n+1})} \\
&\quad + \frac{y_n((1 + \tan(A_n + B_{n+1}))(1 + \tan(A_{n+1} + B_{n+1})) - I_n)}{1 + \tan(A_{n+1} + B_{n+1})} \\
&= \frac{(\delta_n + y_n)(1 + \tan(A_n + B_{n+1}))(1 + \tan(A_{n+1} + B_{n+1}))}{1 + \tan(A_{n+1} + B_{n+1})} \\
&\quad + \frac{-\delta_n(1 + \tan(A_n + B_{n+1})) - y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\
&= (\delta_n + y_n)(1 + \tan(A_n + B_{n+1})) \\
&\quad + \frac{-\delta_n(1 + \tan(A_n + B_{n+1})) - y_n I_n}{1 + \tan(A_{n+1} + B_{n+1})} \\
&= c_n + (\delta_n - \delta'_n),
\end{aligned}$$

where in the last equality we use the definitions of c_n (2.13) and δ'_n (2.16).

(2) Recall,

$$\begin{aligned}
\Delta_n^\varepsilon &= \{(y_1, y_2) : y_1 = \varepsilon, (\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 < \\
&\quad < y_2 < (\varepsilon + x_n) \cdot \tan(A_n + B_{n+1}) - 1\}.
\end{aligned}$$

We start by proving the inequality

$$(\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 \leq c_n,$$

for some $\delta'_n < \varepsilon < \delta_n$.

Let $\delta'_n < \varepsilon < \delta_n$. Then the inequality

$$(\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 \leq (\delta_n + 1) \tan(A_n + B_n) + 1$$

holds. Therefore, the definition of P'_n (2.9) yields

$$\begin{aligned} & (\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 \\ & \leq \delta_n \cdot \tan(A_n + B_{n+1}) + (1 + \tan(A_n + B_n) - \delta_n P'_n). \end{aligned}$$

The definition of y_n (2.12) implies

$$(\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 \leq (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n.$$

Eventually, the definition of c_n (2.13) implies

$$(\varepsilon + 1) \cdot \tan(A_n + B_n) + 1 \leq c_n,$$

and hence the inequality.

In order to prove the inequality

$$c_n + (\delta_n - \delta'_n) \leq (\varepsilon + x_n) \cdot \tan(A_n + B_{n+1}) - 1,$$

it is enough to show it for $\varepsilon = \delta'_n$.

We also use two relations:

$$(4.10) \quad \delta_n - \delta'_n = \delta'_n \cdot \tan(A_{n+1} + B_{n+1}) - \delta_n \cdot \tan(A_n + B_{n+1}) - y_n \cdot I_n,$$

$$(4.11) \quad y_n = \frac{\delta_n P_n - x_n \tan(A_n + B_{n+1}) + 1}{1 + \tan(A_{n+1} + B_{n+1})}.$$

Notice that these statements follow directly from the definition of δ'_n (2.16) and the equality

$$d_n = (\delta_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1$$

from the Lemma 4.4, respectively.

Due to

$$\delta'_n < \delta_n,$$

and the definition of P_n (2.8), the inequality

$$\begin{aligned} & -\delta_n P_n + x_n \cdot \tan(A_n + B_{n+1}) - 1 + \delta'_n \cdot \tan(A_{n+1} + B_{n+1}) \\ & \leq (\delta'_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1 \end{aligned}$$

holds. Therefore, the equality (4.11) implies

$$\begin{aligned} & \delta'_n \cdot \tan(A_{n+1} + B_{n+1}) - y_n \cdot (1 + \tan(A_{n+1} + B_{n+1})) \\ & \leq (\delta'_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1. \end{aligned}$$

From here, using (4.10), the inequality

$$\begin{aligned} & (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n + (\delta_n - \delta'_n) \\ & \leq (\delta'_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1 \end{aligned}$$

follows. The definition of c_n (2.13) yields,

$$\begin{aligned} & (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n + (\delta_n - \delta'_n) \\ & \leq (\delta'_n + x_n) \cdot \tan(A_n + B_{n+1}) - 1, \end{aligned}$$

and the inequality is proved. $\square$

Lemma 4.6. *Let $I_n, P_n, P'_n, \delta_n, x_n, y_n, c_n, d_n, U_n, \delta'_n, S_n = (x_{S_n}, y_{S_n}), r_n, y_{P_1}$ and y_{P_2} be as in the Proposition 2.3. Then*

- (1) $y_{P_1} \leq c_n$,
- (2) $c_n + (\delta_n - \delta'_n) \leq y_{P_2}$.

Proof. (1) The inequality

$$\varepsilon < \delta_n,$$

implies

$$(\varepsilon + y_n) \cdot \tan(A_n + B_{n+1}) + y_n \leq (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n.$$

Therefore, together with (2.21) and (2.13), the inequality

$$y_{P_1} \leq c_n$$

follows.

- (2) To prove the second inequality, we use $\delta'_n < \varepsilon$ and (4.10).

Namely,

$$\delta'_n \leq \varepsilon$$

implies

$$\begin{aligned} & \delta'_n \tan(A_{n+1} + B_{n+1}) - y_n(1 + \tan(A_{n+1} + B_{n+1})) \\ & \leq \varepsilon \cdot \tan(A_{n+1} + B_{n+1}) - y_n(1 + \tan(A_{n+1} + B_{n+1})). \end{aligned}$$

From here we obtain

$$\begin{aligned} & (\delta_n + y_n) \cdot \tan(A_n + B_{n+1}) + y_n + \\ & + \delta'_n \cdot \tan(A_{n+1} + B_{n+1}) - \delta_n \cdot \tan(A_n + B_{n+1}) - y_n \cdot I_n \leq \\ & \leq (\varepsilon - y_n) \cdot \tan(A_{n+1} + B_{n+1}) - y_n \end{aligned}$$

Using (2.22), (2.13) and (4.10) it follows

$$c_n + (\delta_n - \delta'_n) \leq y_{P_2},$$

and hence the Lemma. □

Lemma 4.7. *Let $I_n, P_n, P'_n, \delta_n, y_n, b_n$ and M be as in the proof of Theorem 2.14, i.e. from Proposition 2.2. Then:*

$$\limsup_{n \rightarrow \infty} \frac{M}{y_n} < \infty.$$

Proof. By the definition of δ_n (2.10), it follows:

$$\begin{aligned} y_n &= \frac{1 + \tan(A_n + B_n) - \delta_n \cdot P'_n}{1 + \tan(A_n + B_{n+1})} \\ (4.12) \quad &= \frac{P_n^2 \cdot (1 + \tan(A_n + B_n))}{H_n}, \end{aligned}$$

where H_n is like in the proof of the Proposition 2.2.

From here and (2.10) it follows

$$(4.13) \quad \frac{\delta_n}{y_n} = \frac{I_n \cdot (1 + \tan(A_{n+1} + B_{n+1}))}{P_n^2}.$$

Also, by (4.12) and the definitions of δ_n (2.10) and x_n (2.11), it follows

$$(4.14) \quad \frac{x_n}{y_n} = R_1 + R_2 + R_3,$$

where

$$\begin{aligned} R_1 &= 2 \cdot \frac{1 + \tan(A_n + B_{n+1})}{\tan(A_n + B_{n+1})} \cdot \frac{1 + \tan(A_{n+1} + B_{n+1})}{P_n}, \\ R_2 &= \frac{1 + \tan(A_n + B_{n+1})}{(1 + \tan(A_n + B_n)) \cdot \tan(A_n + B_{n+1})}, \\ R_3 &= \frac{I_n \cdot (1 + \tan(A_{n+1} + B_{n+1}))}{P_n^2} \frac{P'_n}{\tan(A_n + B_{n+1})} \frac{1}{1 + \tan(A_n + B_n)}. \end{aligned}$$

Combining (4.13) and (4.14), the conditions (2.3) and (2.4) imply

$$\limsup_{n \rightarrow \infty} \frac{M}{y_n} < \infty,$$

and hence the proposition. $\square$

REFERENCES

1. K. Bogdan, *The boundary Harnack principle for the fractional Laplacian*, Studia Math. 123 (1997), no.1, 43–80. MR 1438304 (98g:31005)
2. K. Bogdan and P. Sztonyk, *Harnack's inequality for stable Lévy processes*, Potential Anal. 22 (2005), no.2, 133–150. MR 2137058 (2005m:31015)
3. K. Bogdan and P. Sztonyk, *Estimates of the potential kernel and Harnack's inequality for the anisotropic fractional Laplacian*, Studia Math. 181 (2007), no.2, 101–123. MR 2320691 (2008k:31015)
4. Z.-Q. Chen, *Lecture Notes: Potential Theory for Jump Processes; Summer School 2012 on Nonlocal Operators*, 2012.
5. N. Ikeda and S. Watanabe, *On some relations between the harmonic measure and the Lévy measure for a certain class of Markov processes*, J. Math. Kyoto Univ. 2 (1962), 79–95. MR 0142153 (25 #5546)
6. T. Jakubowski, *On Harnack inequality for α -stable Ornstein-Uhlenbeck processes*, Math. Z. 258 (2008), no. 3, 609–628. MR 2369047 (2009b:60242)
7. N. S. Landkof, *Foundations of modern potential theory*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen mit besonderer Berücksichtigung der Anwendungsgebiete ; 180, vol. 180, Springer, Berlin [u.a.], 1972.
8. E. Priola, *Pathwise uniqueness for singular SDEs driven by stable processes*, Osaka J. Math. 49 (2012), no. 2, 421–447. MR 2945756
9. K. Sato, *Lévy processes and infinitely divisible distributions*, digital print. ed., Cambridge studies in advanced mathematics; 68, vol. 68, Cambridge Univ. Press, Cambridge [u.a.], 2007 (eng).
10. P. Sztonyk, *Estimates of tempered stable densities*, J. Theoret. Probab. 23 (2010), no. 1, 127–147. MR 2591907 (2011b:60183)
11. S. J. Taylor, *Sample path properties of a transient stable process*, J. Math. Mech. 16 (1967), 1229–1246. MR 0208684 (34 #8493)

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, UNIVERSITÄTSSTRASSE 25, 33615 BIELEFELD

E-mail address: msertic@math.uni-bielefeld.de